

Today:

Clouds from instability, orographics and weather systems

**Admin stuff:**

- Idea Forge resources: online in Course info.

Also, your students can reserve Area 51 (it's a conference room on the ground level, no windows) for a dark area. The calendar for the room is available here:

[https://calendar.google.com/calendar/embed?](https://calendar.google.com/calendar/embed?src=rb3ml0g7q00aujok9kcn3bn9ms@group.calendar.google.com&ctz=America/Denver&mode=week)

[src=rb3ml0g7q00aujok9kcn3bn9ms@group.calendar.google.com&ctz=America/Denver&mode=week](https://calendar.google.com/calendar/embed?src=rb3ml0g7q00aujok9kcn3bn9ms@group.calendar.google.com&ctz=America/Denver&mode=week)

Email [Rebecca.komarek@colorado.edu](mailto:Rebecca.komarek@colorado.edu) or Lauren Wheeler

([lauren.wheeler@colorado.edu](mailto:lauren.wheeler@colorado.edu)) to reserve the space. Pick up a key from Rebecca (Becky), the machine shop office, or the front desk, near out east entrance.

- <http://matadornetwork.com/bnt/60-insane-cloud-formations-from-around-the-world-pics/> From Michael Lloyd
- FV 2003 alumnus Emrys Hall shot this cloud image yesterday: He works at NOAA, measuring water vapor, sending up special weather balloons  
<http://www.esrl.noaa.gov/gmd/ozwv/wvap/>



- Please edit your Get Wet post to include your report. Do this for all your reports this semester; add them to your image/vid post on the report due date.

- Cloud image submission: Include
  - 1) your edited image
  - 2) your original (unedited) image
  - 3) the appropriate Skew-T diagram
  - 4) a short statement of cloud type and stable or unstable atm.
  - 5) Post on Flowvis.org. Edit your post date to match your cloud date and time.


---

### Clouds = droplets or ice MOVING UPWARDS



Lift mechanisms:


1. Instability: creates Cumulus clouds
2. Orographics: terrain, mountains
3. Synoptic scale weather systems; local instability. Both at warm and cold fronts; cold air pushes under in a cold front, warm air overruns in a warm front.
4. Convergence: shoreline temperature differences and cyclonic uplift

# 1. Instability driven clouds



If atmosphere is UNSTABLE, the heated air will continue to go up!



  
Dark ground (plowed field etc.) can create

<http://www.k3jae.com/wxstormdevelopment.php>

local hot spot, starting a thermal. Mountain uplift can also trigger start of cycle.


Stratocumulus  
Formation mechanisms:

<http://www.flowvis.org/category/flow-categories/clouds/stratocumulus/>



<http://www.flowvis.org/2013/04/11/stratocumulus-boulder-co-18th-of-february-2013-at-1131-a-m/>

- 1) Cumulus joined together, caused by an inversion, a stable layer that stops upward convection
- 2) Stratus broken up. Top reflects UV, visible light, cools (maybe radiates IR to space). Bottom absorbs IR from the earth, warms. Cool on top, warm on the bottom = unstable, wants to turn over, breaking up stratus layer. Stratocumulus stratiformis



Partial rule of thumb  
Cumulus = from instability; local uplift  
Stratus = more stable, from widespread uplift

These are GENUS

For info on Species, Varieties and Accessory Clouds, see  
<http://cloudappreciationsociety.org/collecting/about-cloud-classifications/>

Interesting book on how clouds were first classified and named ~1804, by Luke Howard  
Richard Hamblyn, *The Invention of Clouds: How an Amateur Meteorologist Forged the Language of the Skies* (Picador, 2002).

## 2: Orographic clouds, caused by topography, i.e. mountains

Most common interesting cloud in spring is the

standing  
Alto cumulus lenticularis (higher than 6500 ft above local ground level) ACSL  
or  
Stratocumulus lenticularis (lower)  
or  
Mountain Wave Cloud, trapped or lee


requires STABLE atmosphere: note exception to unstable/cumulus pairing

STANDING WAVE  
Clouds Produced by Vertically **Trapped** Mountain Waves

Thomas Carney et al.,  
*AC 00-57 Hazardous Mountain Winds and Their Visual Indicators*  
(Federal Aviation Administration, 1997),  
<http://rgl.faa.gov/Regu>




Mountain Waves and  
Their Visual Indicators  
(Federal Aviation  
Administration, 1997),  
[http://rgl.faa.gov/Regulatory\\_and\\_Guidance\\_Library/rgAdvisoryCircular.nsf/0/780437D88CBDAFD086256A94006FD5B8?OpenDocument](http://rgl.faa.gov/Regulatory_and_Guidance_Library/rgAdvisoryCircular.nsf/0/780437D88CBDAFD086256A94006FD5B8?OpenDocument).



Clouds that sit right  
on the Divide =  
FOEHN cloud wall.  
From air being forced up over  
the mountains

Altostratus lenticularis. Typically 1 to 5 wave crests.

Clouds stay stationary, but may move off and reform periodically



Ben Britton, FV 2010

If there's more wave crests, or short wavelengths, it's probably NOT a mountain wave cloud; more likely altostratus undulatus, from gravity waves in the atmosphere, like ripples on a liquid surface.

<http://www.colorado.edu/MCEN/flowvis/galleries/2007/assignment2.html>





Tracy Eliasson FV 2007

Could also be from wind shear, via the Kelvin Helmholtz instability



Rare to be able to see cross section like this


<http://cloudappreciationsociety.org/collecting/terry-robinson/>



Minute paper: Which way is the wind going?  
Where is it faster?



Colin Stewart FV 2012 Clouds 1



FÖHN

ADABATIC

Foehn clouds suggest winds coming over the mountains: the presence of a CHINOOK (pre-cold-front, warm, strong, downslope winds, or a BORA (post-cold-front, cold, strong, downslope winds). Also called cap clouds.


KATABATIC

3: Synoptic uplift = weather system clouds.

Weather system progressions; 'synoptic scale' uplifts (1000 km across). Any type of cloud is possible.

Inserted from: -file:///C:/Users/hertzber/Documents/01CLASSES/FlowVis/Content/scanned\_images/TypWeatherSystem.tif-



The Cloudspotter's Guide pg186 THE HIGH CLOUDS



How cloud formations can develop as a region of low pressure, or 'depression', passes over. Those who think this looks complicated will be depressed to learn that it is in fact a very simplified diagram of a weather system.


Pressure ↑


XXXXXXXX



Low Pressure System: Air tries to move into low.  
 — Coriolis makes it turn left = counterclockwise circulation. Typically unstable. **CYCLONIC**

High pressure system: Air tries to move out.  
 Coriolis makes it turn right = clockwise circulation. Weak or nonexistent fronts, so no instability. **ANTICYCLONIC**






[http://earth.usc.edu  
/~stott/Catalina/WeatherPatterns.html](http://earth.usc.edu/~stott/Catalina/WeatherPatterns.html)

Divergence aloft creates convergence and lift at surface. Pumping action.

#### 4: Convergence uplift along shorelines



CloudClassificationTable.pdf; Copyrighted, but available in D2L.

Also see

[Cloud types for observers \(PDF, 4 MB\) - Met Office 45 pgs](#)

The Cloud Spotter's Guide  
CLOUD CLASSIFICATION TABLE  
Gayin Pretor-Pinney, Berigee Press 2006  
Clouds are classified according to a Latin 'Linnaean' system (similar to the one used for plants and animals), which is based on their heights and appearance. Most clouds fall into one of ten basic groups, known as 'genera'. They can further be defined as one of the possible 'species' for that genus, and any combination of the possible 'varieties'. There are also various accessory clouds and supplementary features that sometimes appear in conjunction with the main cloud types. (If all this Latin freaks you out, don't worry - it freaks me out too.)

GENUS	SPECIES (CAN ONLY BE ONE)	VARIETIES (CAN BE NONE OR MORE THAN ONE)	ACCESSORY CLOUDS AND SUPPLEMENTARY FEATURES
Cumulus	humilis		pileus
	mediocris	radiatus	velum
	congestus		virga
	fractus		praecipitatio
Cumulonimbus (extends through all three levels)	calvus		praecipitatio
	capillatus	(none)	virga
			velum
			pernis
		arcus	
		stercus	
		tuba	
	nebulosus	opacus	mamma



The Cloud Spotter's Guide

CLOUD CLASSIFICATION TABLE

Gaylin Pretor-Painey, Berigee Press 2006

Clouds are classified according to 4 Latin "linear" stems (similar to the one used for plants and animals), which is based on their heights and appearance. Most clouds fall into one of ten basic groups, known as "genera". They can further be defined as one of the possible "species" for that genus, and any combination of the possible "varieties". There are also various accessory clouds and supplementary features that sometimes appear in conjunction with the main cloud types. (If all this Latin freaks you out, don't worry - it freaks me out too.)

GENUS	SPECIES (CAN ONLY BE ONE)	VARIETIES (CAN BE NONE OR MANY)	ACCESSORY CLOUDS AND SUPPLEMENTARY FEATURES
Cumulus	humilis		velum parvus
	mediocris	radiatus	virga tuba
	congestus		praecipitatio
	fractus		praecipitatio pileus
Cumulonimbus (extends through all five levels)	calvus		virga velum
	capillatus	(none)	parvus arcus tuba
Stratus	nebulosus	opacus	
	fractus	transversus undulatus	praecipitatio
		transversus perfoliatus	
Stratocumulus	stratiformis	opacus	mammata
	lenticularis	duplicatus	virga
	castellatus	undulatus	praecipitatio
		radiatus lacunosus	
Altostratus	stratiformis	perfoliatus	
	lenticularis	opacus	virga
	castellatus	duplicatus	mammata
	floccus	undulatus	
		radiatus lacunosus	
Altostratus	transversus	opacus	virga praecipitatio
	(none)	duplicatus	parvus mammata
		undulatus radiatus	
			praecipitatio
Nimbostratus (extends through most above five levels)	(none)	(none)	virga parvus
Cirrus	fibrosus	intortus	
	unicus	radiatus	
	spinosus	verticillatus	mammata
	castellatus	duplicatus	
	floccus		
	stratiformis	undulatus	virga
Circumcunulus	lenticularis	undulatus	virga mammata
	castellatus	lacunosus	
	floccus		
Cirrostratus	fibrosus	duplicatus	(none)
	nebulosus	undulatus	

HOW TO SPOT  
**CUMULUS CLOUDS**

**C**umulus are low, detached, puffy clouds that develop vertically in rising mounds, domes or towers, and have generally flat bases. Their upper parts often resemble cauliflower and they appear brilliant white when reflecting high sunlight, but can look dark when the sun is behind them. Cumulus tend to be randomly scattered across the sky.

**TYPICAL ALTITUDES\*:**  
 2,000-3,000ft  
**WHERE THEY FORM:**  
 Worldwide, except in Antarctica (the ground is too cold for thermals).  
**PRECIPITATION (REACHING GROUND):** Generally none, except for brief showers from congestus.



Cumulus humilis



Cumulus mediocris



Cumulus congestus

**CUMULUS SPECIES:**  
**HUMILIS:** Minimal vertical extent. They look flattened and appear wider than they are tall. Do not cause precipitation.  
**MEIOCRIS:** Moderate vertical extent. Might show protuberances and sproutings at the top. Appear as tall as they are wide. Do not cause precipitation.  
**CONGESTUS:** Maximum vertical extent. The tops are like cauliflowers. Appear taller than they are wide. Cause brief downpours.  
**FRACTUS:** Ragged edges and broken up. Can form in the moist air below rain clouds.

**CUMULUS VARIETIES:**  
**RADIATUS:** When Cumulus have formed into rows, or 'cloud streets', which are roughly parallel to the wind direction. Due to perspective, the rows appear to converge towards the horizon.



Cumulus radiocirrus radiatus

**NOT TO BE CONFUSED WITH...**  
**STRATOCUMULUS:** Cumulus clouds are detached, not joined into a layer like Stratocumulus.  
**ALTOCUMULUS:** Cumulus are not usually as regularly spaced as a layer of the higher AltoCumulus. The clouds also look larger than the clumps of the AltoCumulus. When they are above the cloudspotter, Cumulus appear larger than the width of three fingers, held at arm's length.  
**CUMULONIMBUS:** which often develops from a large Cumulus congestus. A cloud is still a Cumulus when its upper region has a sharp outline, compared with the softer top of the Cumulonimbus.


\* These approximate altitudes (above the surface) are for mid-latitude regions.

Top right: Michael Smith (twitter: 129) Bottom: Paul Cooper (twitter: 1243)

**HOW TO SPOT  
CUMULONIMBUS CLOUDS**

**C**umulonimbus are thunderstorm clouds, characterised by their enormous height. They are typically tall enough to reach the top of the troposphere, where they spread out in plumes of ice particles that can appear smooth, fibrous or striated. They have dark bases and produce heavy showers – often of hail – which can be accompanied by thunder and lightning.

**TYPICAL ALTITUDES\*:** 2,000–45,000ft  
**WHERE THEY FORM:** Common in tropical and temperate regions. Rare in polar ones.  
**PRECIPITATION (REACHING GROUND):** Heavy downpours, often of hail.



**CUMULONIMBUS SPECIES:**

The two species are distinguished by the appearance of the cloud's top.  
**CALVUS:** When the upper region is of soft indistinct flattened mounds, without any fibrous or striated appearance.  
**CAPILLATUS:** When the upper region is cirrus-like and fibrous or striated, often in the shape of an anvil, plume or a disorderly mass of white hair.

**CUMULONIMBUS VARIETIES:**

There are no official varieties.



\* These approximate altitudes (above the surface) are for mid-latitude regions.

**NOT TO BE CONFUSED WITH...**

**NIROBORATUS:** which is a dark, ragged precipitating layer, covering the sky. It can look similar to a Cumulonimbus that is directly overhead (and also appears to cover much of the sky) but the precipitation will tend to be more steady and more persistent than the short heavy showers of the Cumulonimbus. If thunder, lightning or hail is present, then the cloud is a Cumulonimbus.  
**CUMULUS CONGESTUS:** from which a Cumulonimbus often develops. Seen from a distance, the cloud is said to have changed into a Cumulonimbus when parts of its upper region begin to lose their sharp edges, due to the droplets freezing into ice crystals. Thunder, lightning or hail will also identify the Cumulonimbus.

Top left: Bob Marshall (Creative Commons); Top right: Mike Davis (Creative Commons); Bottom: Bob Marshall (Creative Commons)

I  
L  
L  
U  
S  
T  
R  
A  
T  
I  
O  
N  
S  
B  
Y  
C  
H

**HOW TO SPOT  
STRATUS CLOUDS**

**S**tratus are grey layers or patches of cloud, with very diffuse edges. They are the lowest forming of all the cloud genera, sometimes appearing at ground level, when they are called fog or mist.

**TYPICAL ALTITUDES\*:** 0–6,500ft  
**WHERE THEY FORM:** Worldwide. Most commonly around coasts and mountainous.  
**PRECIPITATION (REACHING GROUND):** No more than occasional drizzle, snow or snow grains.



Stratus stratiformis tenuis



Stratus

**NOT TO BE CONFUSED WITH...**

**CONGESTUS:** which is a high layer cloud that can look similar to a very thin Stratus. Being made of ice, however, it has a white top.

**ALTOFRATUS:** which is a mid-level layer cloud, often consisting of droplets, like Stratus. Through a layer of Stratus, the outline of the Sun (even if it is dimmed) is less diffuse, compared with the 'ground-glass' appearance of the Altostratus.

**NIMBIFRATUS:** which is a thick, dark layer of precipitating cloud that might be confused with a thick Stratus. But this has a less ragged base than the Nimbifratus and produces lighter precipitation.

**STRATUS SPECIES:**

**NIROBORATUS:** By far the most common, when it is in a grey, generally featureless layer.

**FRACTUS:** When it is in separate, ragged sheets of grey cloud. This can appear in the region below precipitating clouds, when it is called 'pannae'. Though not particularly thick, these sheets can look quite dark against the base of the cloud above.

**STRATUS VARIETIES:**

**OPACUS:** When the layer is thick enough to completely mask the sun or moon.

**FRANULOSUS:** When it is thin enough to show the outline of the sun or moon.

**NIROBORATUS:** A rare variety, in which the layer has more like undulations to its surface. The surface of Stratus is rarely distinct enough for this to be observed.


\* These approximate altitudes (above the surface) are for mid-latitude regions.

P  
f  
  
M  
w  
'c  
't  
N  
  
G  
E  
H  
E  
f  
n  
N  
  
ir  
A  
b  
B  
n  
M  
  
C  
n  
n  
O  
H  
M  
M  
C  
  
n  
M  
w

HOW TO SPOT  
**STRATOCUMULUS CLOUDS**

**S**tratocumulus are low layers or patches of cloud, with well-defined bases. They are usually composed of clumps or rolls, and often show strong variations in tone - from bright white to dark grey. Their cloud elements may be joined into continuous, unbroken layers or have gaps between them.

**TYPICAL ALTITUDES\*:**  
2,000-4,500'  
**WHERE THEY FORM:**  
Worldwide - it's a very common cloud.  
**PRECIPITATION**  
**(REACHING GROUND):**  
Occasionally light rain, snow or snow pellets.



Stratocumulus stratiformis opacus...



...and perlucidus

**STRATOCUMULUS SPECIES:**  
**STRATIFORMIS:** The most common, when the clumps or rolls extend over a large area. A 'roll cloud' is a particular formation, in the shape of a large, individual tube of cloud.  
**LENTICULARIS:** When one or more mass of cloud is in a smooth, solid-looking almond or lens shape.  
**CASTELLANUS:** When the elements have crenellated tops.

**STRATOCUMULUS VARIETIES:**  
**OPACUS:** When the layer is thick enough to completely mask the sun or moon.  
**TRANSLUCIDUS:** When it is thin enough to show the outline of the sun or moon.  
**PERLUCIDUS:** When there are gaps between the cloud elements.

**DUPPLICATUS:** When there are layers at different altitudes, sometimes partly merged.  
**UNDULATUS:** When the elements are arranged in nearly parallel lines.  
**RADIATUS:** When lines of closely bunched elements appear to converge towards the horizon.  
**LACUNOSUS:** When the layer shows large net-like holes fringed with cloud.

**NOTE TO BE CONSIDERED WITH:**  
**CUMULUS:** which is also clumpy, well defined, and forms at similar altitudes. The elements of Stratocumulus tend to be closer together and to have flatter tops.  
**ALTOCUMULUS:** which is a mid-level layer of cloudlets. These appear smaller than the Stratocumulus elements, which - looking above 30' from the horizon - appear larger than the width of three fingers, held at arm's length.  
**STRATUS:** which is a low, indistinct layer, with much less variation in tone and less definition than Stratocumulus.

\* These approximate altitudes (above the surface) are for mid-latitude regions

on  
tog  
aw  
po  
rat  
fu  
ab  
esc  
th

cle  
esi  
Ni  
fer  
co  
so

pe  
th  
dr  
wi  
wo  
G  
A

cl  
uj

S  
b  
cl  
ai  
g  
it  
te

HOW TO SPOT  
**ALTOCUMULUS CLOUDS**

**A**ltocumulus are mid-level layers or patches of cloudlets, in the shape of rounded clumps, rolls or almonds/lenses. These are white or grey, and the sides away from the Sun are shaded. Altocumulus are usually composed of droplets, but may also contain ice crystals.

**TYPICAL ALTITUDES\***:  
 6,500–18,000ft  
**WHERE THEY FORM**:  
 Worldwide.  
**PRECIPITATION**  
 (REACHING GROUND):  
 Very occasionally causes light rain.

**ALTOCUMULUS SPECIES:**

**STRATIFORMIS**: Most common, when the cloudlets extend over a large area.


**LENTICULARIS**: When it is in the form of one or more individual almond- or lens-shaped masses that appear dense, with pronounced shading.

**CASTELLANUS**: When the cloudlets have crenellated tops.

**FLOCCUS**: When the cloudlets are Cumulus-like tufts, with ragged bases, often with fibrous trails (virga) of ice crystals falling below.



*Altocumulus stratiformis undulata*



*Altocumulus lenticularis*

*Altocumulus floccus*

**ALTOCUMULUS VARIETIES:**

**OPACUS**: When the layer is thick enough to completely mask the sun or moon.

**TRANSALCINUS**: When it is thin enough to show the outline of the sun or moon.

**FRILLOSCINUS**: When there are gaps between the cloudlets.

**DUPLICATUS**: When there are layers at different altitudes, sometimes partly merged.

**UNDULATUS**: When the cloudlets are arranged in nearly parallel lines.

**RADIATUS**: When long lines of them appear to converge towards the horizon.

**LACUNOSUS**: When the layer shows net-like holes fringed with cloud.

**NOT TO BE CONFUSED WITH:**

**CIRROCUMULUS**: which is a higher layer of cloudlets, that appear like little grains of salt. Looking above 30° from the horizon, the larger Altocumulus cloudlets generally appear the width of between one and three fingers, held at arm's length.

Also, these exhibit shading, which those of Cirrocumulus don't.

**CIRREUS**: which is a high cloud, whose streaks of falling ice crystals can resemble Altocumulus cloudlets showing virga, but do not have their dense-looking heads.

**ALTOCUMULUS**: which is a mid-level cloud, whose streaks of falling ice crystals can resemble Altocumulus cloudlets showing virga, but do not have their dense-looking heads.

\* These approximate altitudes (above the surface) are for mid-latitude regions.

Left: Stephen Cook (number 122). Below left: Mike Cook (number 149).  
 Below right: Terry Hahn (number 122).

la  
 w  
 b  
 c  
 d  
 t  
 n  
 o  
 f  
 'j  
 h  
 t

(  
 r  
 v  
 c  
 c  
 t  
 i  
 i  
 l

## HOW TO SPOT ALTOSTRATUS CLOUDS

**A**ltostratus are mid-level layers of grey cloud, which are either featureless or fibrous in appearance, and typically extend over an area of several thousand square miles. Usually composed of both water droplets and ice crystals, they are often thin enough in parts to reveal the position of the sun, which appears as if through ground glass. Altostratus can cause a white or (when very thin) coloured 'corona' (disc of light) around the sun or moon.




*Altostratus nimbocubus*

**TYPICAL ALTITUDES\*:**  
6,500-23,000ft

**WHERE THEY FORM:**  
Worldwide. More common in the middle latitudes.

**PRECIPITATION (REACHING GROUND):** Usually not, but occasionally light rain or snow.

**ALTOSTRATUS SPECIES:**  
There are no species, as the cloud's appearance is so uniform.



*Altostratus radiatus*

### ALTOSTRATUS VARIETIES:

**OPACUS:** When the cloud layer is generally thick enough to mask the position of the sun or moon.

**FRANKLICOIDES:** When it is generally thin enough to show the position of the sun or moon.

**DUPLICATUS:** When there is more than one layer at different altitudes, these often being partly merged. This is generally only visible when, by the light of a low sun, the higher layer is lit and the lower is in shadow, or when shearing winds cause the striations of the layers to differ.

**UNDULATUS:** When the layer shows largely parallel undulations.

**RAGNATUS:** When lumpy undulations appear to converge toward the horizon.

### NOT TO BE CONFUSED WITH...

**CIRROSTRATUS:** which is a higher layer of ice crystals that looks like a thin, milky veil across the sky, and often thickens and lowers to develop into Altostratus. The Altostratus will tend to be more opaque, making the sunlight too diffuse for objects to cast shadows, as they do below Cirrostratus. White coloured or white discs of light, called coronae, can appear around the sun/moon through Altostratus, this cloud will not cause the 'halo phenomena' of the Cirrostratus.

**NIMBOSTRATUS:** which is a thick, dark layer of precipitating cloud that often develops out of an Altostratus. Generally darker, it produces considerably heavier rain or snow.

\* These approximate altitudes (above the surface) are for mid-latitude regions.

From: *www.zimbardo.com* (member 18/11/10) *Flight - October Skyline* (number 1175)

HOW TO SPOT  
**NIMBOSTRATUS CLOUDS**

**N**imbostratus are thick, grey, featureless layers of cloud that cause prolonged, continuous, often heavy, rain, snow or ice pellets. They tend to have very diffuse bases, as a result of all the falling precipitation. Nimbostratus are the deepest of all the layer clouds - sometimes extending from 2,000ft up to around 18,000ft - and generally extend over many thousand square miles. As with other precipitating clouds, the falling precipitation can cause Stratus fractus to form in the air below Nimbostratus clouds. These are known as 'pannus' and appear as shreds of cloud, looking darker than the underside of the Nimbostratus. When these join together, they tend to lower the bases of Nimbostratus clouds even further. They are invariably thick enough to completely hide the sun or moon.

**NOT TO BE CONFUSED WITH:**  
**ALTOSTRATUS:** which is a thinner - though also indistinct - layer of cloud. Nimbostratus is always darker than it and, by definition, produces precipitation. Altostratus only does sometimes, and this will generally be light. Whilst the position of

**TYPICAL ALTITUDES\*:**  
 2,000-18,000ft

**WHERE THEY FORM:**  
 Worldwide. More common in middle latitudes.

**PRECIPITATION (REACHING GROUND):** Causes moderate to heavy rain or snow (steady and prolonged).

**NIMBOSTRATUS SPECIES:**  
 There are no species, as the cloud's appearance is so uniform.

**NIMBOSTRATUS VARIETIES:**  
 There are no varieties, as the cloud's appearance is so uniform.

the sun can generally be determined through at least part of a layer of Altostratus, it will never be so through a Nimbostratus.

**CUMULONIMBUS:** which, when observed from directly below, can also appear as a very dark layer, covering the whole sky. The precipitation falling from a Nimbostratus will not generally be as heavy and will be more prolonged and continuous, compared with the sudden showers of the Cumulonimbus. Nor will the Nimbostratus produce its hail, thunder or lightning.



Nimbostratus - never a pretty sight

\* These approximate altitudes (above the surface) are for mid-latitude regions.

David Evans (number 117)

## HOW TO SPOT CIRRUS CLOUDS

**C**irrus are the highest of the ten main cloud types. In the form of delicate, white streaks, patches or bands of falling ice crystals, they are detached from each other, and have fibrous or silky appearances. Cirrus rarely appear very thick. They are often seen with the other high clouds, Cirrostratus and Cirrocumulus and, like them, can show 'halo phenomena' around the sun or moon.



Cirrus uncinus



Cirrus floccus



Cirrus vertebratus

### CIRRUS SPECIES:

**FIBRATUS:** When it is in the form of straight or curved filaments that are mostly distinct from each other and do not terminate in hooks or clumps.

**UNCINUS:** When its 'fallstreaks' are the shape of hooks or comas.

**SPISSATUS:** The thickest Cirrus - when it is in patches that appear grey in front of the sun - which tends to originate from the arrival of a Cumulonimbus.

**CASTELLATUS:** When it is in the form of small distinct clumps with crenellated tops.

**FLOCCUS:** When it is in the form of independent small round tufts, which often show trails of ice crystals falling from them.

### CIRRUS VARIETIES:

**INTORTUS:** When the fallstreaks are irregular and tangled.

**RADIATUS:** When the filaments are in parallel bands, usually aligned to the wind at high altitude, which converge towards the horizon, due to perspective.

### NOT TO BE

**CONFUSED WITH...**

**CIRROSTRATUS:** which looks like a thin, milky smooth or fibrous veil across the sky. Cirrus, by contrast, is in separated streaks, fibres or patches.

**CIRROCUMULUS:** which is a high layer of cloudlets, like grains of salt. Cirrus does not show this finely dappled texture.

### VERTebratus:

When the filaments look like a fish skeleton.

**DUPLICATUS:** When the filaments, streaks or hooks are arranged at more than one altitude, which can be apparent when the winds cause them to point in different directions.

\* These approximate altitudes (above the surface) are for mid-latitude regions.

**TYPICAL ALTITUDES\*:**  
16,500-45,000ft

**WHERE THEY FORM:**  
Worldwide

**PRECIPITATION (REACHING GROUND):** None.

Left: Matthew Reed (November 11/10); Top right: Graham Tibbo (November 10/2)  
Bottom right: Grant King (November 10/19)

re  
w  
th  
  
cl  
'p  
N  
cr  
sp  
cl  
  
th  
di  
of  
Fc  
pl  
  
cl  
re  
di  
th  
co  
  
tr  
to  
va  
th  
od



HOW TO SPOT  
**CIRROCUMULUS CLOUDS**

**C**irrocumulus are high patches of cloud or layers of tiny cloudlets that appear as white grains. These show no shading, even on the sides away from the sun. These cloudlets are generally regularly spaced, and often arranged in ripples, known as the undulatus variety.

**TYPICAL ALTITUDES\*:**  
16,500-45,000ft  
**WHERE THEY FORM:**  
Worldwide.  
**PRECIPITATION (REACHING GROUND):**  
None.


**CIRROCUMULUS SPECIES:**

**STRATIFORMIS:** When it is in an extensive layer, rather than just a patch. A less common species than for other genera.


**LENTICULARIS:** When it is in the form of one or more independent, well-defined, almond- or lens-shaped masses, which have smooth surfaces and are much larger than the grain-like cloudlets of the other species.

**CARTELLANUS:** When, on careful inspection, its cloudlets have crested tops.

**FLOCCUS:** When, on careful inspection, its cloudlets are Cumulus-like, with ragged bases.



*Cirrocumulus stratiformis*



*Cirrocumulus lacunosus undulatus*

**CIRROCUMULUS VARIETIES:**

**UNDULATUS:** When its cloudlets are in a wave-like arrangement of ripples or broad undulations (or both at the same time).

**LACUNOSUS:** When the layer has holes fringed with cloud, like a net or honeycomb.

**NOT TO BE CONFUSED WITH...**

**CIRRUS AND CIRROSTRATUS:** which are streaks and smooth/fibrous layers of high cloud. Cirrocumulus layers, by contrast, are subdivided into many grain-like cloudlets.

**ALTOCUMULUS:** which is a mid-level layer of larger cloudlets. Looking above 30° from the horizon, the smaller Cirrocumulus cloudlets generally appear less than the width of one finger, held at arm's length.

\* These approximate altitudes (above the surface) are for mid-latitude regions.

Top: Steve A. Hartley; middle: 1983; bottom: Steve A. Hartley; middle: 1983

C  
Inde  
a tra  
the :  
way  
appe  
reco,  
delig  
l  
level  
is nc  
how  
sky  
varie  
strat  
a ter  
that  
is pa  
shap

THI  
high  
as t  
rese  
appe  
\  
dete  
ther  
tem  
dep  
form  
imp  
l  
surfi  
the

**HOW TO SPOT  
CIRROSTRATUS CLOUDS**


**C**irrostratus are largely transparent, milky veils of high cloud that look either smooth or fibrous. They tend to cover large areas of the sky, extending over many thousands of square miles, but are often so subtle as to be missed. They do, however, sometimes produce the white or coloured rings, spots or arcs of light around the sun or moon that are known as 'halo phenomena'.

**TYPICAL ALTITUDES\*:**  
16,500-30,000ft


**WHERE THEY FORM:**  
Worldwide.

**PRECIPITATION:**  
None.


**HALO PHENOMENA:**



*Cirrostratus undulatus*



*Cirrostratus causing a '22' Halo' around the moon*



*Cirrostratus fibrosus causing a 'sun-dog' at the same elevation as the sun*

**CIRROSTRATUS SPECIES:**  
**FIBRATUS:** When the cloud veil has a fine fibrous or striated appearance.  
**NEBULOSUS:** When it shows no variation in tone.

**CIRROSTRATUS VARIETIES:**  
**UNDULATUS:** When the veil has a wave-like appearance.  
**DUPLEXATUS:** When there is more than one layer, at different altitudes. This is generally only visible when, by the light of a low sun, the higher layer is lit up when the lower is in shadow, or when shearing winds cause the strata of each layer to differ.

**NOT TO BE CONFUSED WITH...**  
**ALTOSTRATUS:** which is a mid-level, generally thicker, layer cloud. Besides being thinner, the ice crystals of the Cirrostratus can sometimes produce halo phenomena around the sun or moon. These are far less common in Altostratus, which will generally only produce a corona (a white or coloured disc of light).  
**CIRRUS OR CIRROCUMULUS:** which are streaks and grained/rippled layers of high cloud. Cirrostratus, which often appears in conjunction with them, is a more continuous and diffuse layer.

\* These approximate altitudes (above the surface) are for mid-latitude regions.

Consta  
Rome  
a cross  
eris' - '  
Th  
Const  
of that  
safegu  
instruc  
army r  
as the  
Th  
associ  
it later  
as a di  
centre  
Eu  
tally v  
his bo  
it wil  
comp  
Va  
when  
ice cr  
know  
accou  
comm  
I r  
has r  
shape

THE  
whic  
gener  
form